ELSEVIER

Environmental Pollution 128 (2004) I–III

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 128

Numbers 1–2 2004

Persistant Organic Pollutants

Guest Editors: K.C. Jones, A. Sweetman, D. MacKay

1 Editorial
K.C. JONES, A. SWEETMAN (UK), D. MACKAY (Canada)

3 Primary sources of selected POPs: regional and global scale emission inventories
K. BREIVIK (Norway), R. ALCOCK (UK), Y.-F. LI (Canada), R.E. BAILEY (USA), H. FIEDLER (Switzerland), J.M. PACYNA (Norway)
Knowledge of primary emissions is a prerequisite for understanding and predicting POPs on a regional/global scale.

17 Seasonal and long-term trends in atmospheric PAH concentrations: evidence and implications
K. PREVEDOUROS (UK), E. BRORSTRÖM-LUNDÉN (Sweden), C.J. HALSALL, K.C. JONES, R.G.M. LEE, A.J. SWEETMAN (UK)
Seasonality in PAH air concentrations is principally reflecting changes in primary emissions.

29 Distribution of some organochlorine compounds in pine needles from Central and Northern Europe
A. HELLSTRÖM, H. KYLIN (Sweden), W.M.J. STRACHAN (Canada), S. JENSEN (Sweden)
This work shows the distribution of some organochlorines in pine needles in parts of Europe and also shows that current models of air-plant partitioning of hydrophobic compounds are incorrect in important aspects of the uptake processes of airborne contaminants into plants.

49 Soil–air exchange of organochlorine pesticides in the Southern United States
T.F. BIDLEMAN, A.D. LEONE (Canada)
Old pesticide residues in agricultural soils continue to be emitted into the atmosphere.

59 PCB in soils and estimated soil–air exchange fluxes of selected PCB congeners in the south of Sweden
C. BACKE, I.T. COUSINS, P. LARSSON (Sweden)
Soil–air exchange of PCBs is investigated and modelled across Sweden.

73 Air–sea gas exchange of HCHs and PCBs and enantiomers of α-HCH in the Kattegat Sea region
K.L. SUNDQVIST, H. WINGFORS, E. BRORSTÖM-LUNDÉN, K. WIBERG (Sweden)
Air–water exchange of PCBs and HCHs is studied in the Kattegat Sea and shows to vary seasonally.

85 Evaluation of sequentially-coupled POP fluxes estimated from simultaneous measurements in multiple compartments of an air–water–sediment system
A. PALM, I. COUSINS, Ö. GUSTAFSSON, J. AXELMAN, K. GRUNDER, D. BROMAN, E. BRORSTRÖM-LUNDÉN (Sweden)
The magnitude and direction of vertical fluxes of PCBs and PAHs in a coupled air–water–sediment system were simultaneously measured and compared.

99 Current issues and uncertainties in the measurement and modelling of air–vegetation exchange and within-plant processing of POPs
J.L. BARBER, G.O. THOMAS, G. KERSTIENS, K.C. JONES (UK)
Air–plant exchange and plant processing of POPs are key factors affecting global cycling and foodchain transfer.

139 Evidence for the "grasshopper" effect and fractionation during long-range atmospheric transport of organic contaminants
T. GOUIN, D. MACKAY (Canada), K.C. JONES (UK), T. HARNER (Canada), S.N. MEIJER (UK)
Evidence for global scale fractionation and hopping of POPs is reviewed.

149 Latitudinal and seasonal capacity of the surface oceans as a reservoir of polychlorinated biphenyls
E. JURADO (Spain), R. LOHMANN (UK), S. MEIJER (Spain), K.C. JONES (UK), J. DACHS (Spain)
Model calculations estimate the latitudinal and seasonal storage capacity of the surface oceans for PCBs.

163 Investigating the occurrence of persistent organic pollutants (POPs) in the arctic: their atmospheric behaviour and interaction with the seasonal snow pack
C.J. HALSALL (UK)
The fluxes and fate of POPs in snowpacks are key to their behaviour in polar systems.

177 Long-range transport and global fractionation of POPs: insights from multimedia modeling studies
M. SCHERINGER, M. SALZMANN, M. STROEBE, F. WEGMANN, K. FENNER, K. HUNGERBÜHLER (Switzerland)
Multimedia box models are applied to case studies of the behavior of POPs.

doi: 10.1016/S0269-7491(03)00468-8

189 Prediction of overall persistence and long-range transport potential with multimedia fate models: robustness and sensitivity of results
K. FENNER, M. SCHERINGER, K. HUNGERBÜHLER (Switzerland)
Environmental conditions cause variability in substance behavior which need to be considered in chemical ranking schemes.

205 Indicators for persistence and long-range transport potential as derived from multicompartment chemistry–transport modelling
A. LEIP, G. LAMMEL (Germany)
Multicompartment chemistry–transport models can be used to better assess persistence and long-range atmospheric transport potential.

223 BETR-World: a geographically explicit model of chemical fate: application to transport of α-HCH to the Arctic
L. TOOSE, D.G. WOODFINE (Canada), M. MACLEOD (USA), D. MACKAY, J. GOUIN (Canada)
A geographically explicit multi-compartment model is applied to the transport of α-HCH to the Arctic, showing Europe and the Orient are key sources.

241 Modeling transport and deposition of contaminants to ecosystems of concern: a case study for the Laurentian Great Lakes
M. MACLEOD, D. MACKAY (Canada)
Transfer Efficiency is defined as a model-based measure of sources of POPs to the North American Great Lakes.

251 Modelling the fate of persistent organic pollutants in Europe: parameterisation of a gridded distribution model
K. PREVEDOUROS (UK), M. MACLEOD (USA), K.C. JONES, A.J. SWEETMAN (UK)
A regionalised European box model is described, parameterised and tested for Lindane.

263 Evaluating multimedia/multipathway model intake fraction estimates using POP emission and monitoring data
M. MARGNI, D.W. PENNINGTON, C. AMMAN, O. JOLLIET (Switzerland)
The fraction of PCDD/F Emission which transfers to the human population is assessed for Europe using a novel approach.

279 Numerical evaluation of the PCBs transport over the Northern Hemisphere
A. MALANICHEV, E. MANTSEVA, V. SHATALOV, B. STRUKOV, N. VULYKH (Russia)
Transfer of PCBs to the Arctic is assessed using a model of the whole Northern Hemisphere.

291 Effects of time-averaging climate parameters on predicted multicompartmental fate of pesticides and POPs
G. LAMMEL (Germany)
High temporal resolution multi-media modelling approaches are advocated to assess compound behaviour/ranking.

Number 3 2004

Letter to the Editor

303 Letter to the Editor
M. PANELLA, R. AINA, M. RENNA, A. SANTAGOSTINO, L. PALIN (Italy)

Reply to the Letter to the Editor

305 Reply from authors to Letter to the Editor
D. LUTTINGER, L. WILSON (USA)

Research Papers

307 Copper uptake by *Elsholtzia splendens* and *Silene vulgaris* and assessment of copper phytoavailability in contaminated soils
J. SONG (PR China), F.-J. ZHAO (UK), Y.-M. LUO (PR China), S.P. MCGRATH, H. ZHANG (UK)
Both Elsholtzia splendens and Silene vulgaris exclude Cu, and free Cu ions become more available as soil solution pH increases.

317 Antioxidative responses to arsenic in the arsenic-hyperaccumulator Chinese brake fern (*Pteris vittata* L.)
X. CAO, L.Q. MA (USA), C. TU (USA)
At low levels of arsenic exposure, enzymatic antioxidants are important for arsenic detoxification and accumulation in Chinese brake fern, while non-enzymatic antioxidants were more important at high arsenic exposure.

327 Organochlorine contaminants in seven species of Arctic seabirds from northern Baffin Bay
A.H. BUCKMAN, R.J. NORSTROM, K.A. HOBSON (Canada), N.J. KARNOVSKY (USA), J. DUFFE (Canada), A.T. FISK (USA)
Concentrations of organochlorides in high Arctic seabirds are influenced by trophic level, migration, scavenging and biotransformation.

339 Suitability of altai wildrye (*Elymus angustus*) and slender wheatgrass (*Agropyron trachycaulum*) for initial reclamation of saline composite tailings of oil sands
S. RENAULT, C. QUALIZZA, M. MACKINNON (Canada)
Altai wildrye and slender wheatgrass have potential for use in reclamation of amended composite tailings of oil sands.

351 Long-term changes of acidifying deposition in Finland (1973–2000)
J. VUORENMAA (Finland)
Acidic deposition has declined and recovery has taken place in acidified lakes.

363 Vitellogenin content in European eel (*Anguilla anguilla*) in Flanders, Belgium
B.J. VERSONNEN, G. GOEMANS, C. BELPAIRE, C.R. JANSSEN (Belgium)
European eel, Anguilla anguilla, *did not show enhanced plasma VTG concentrations, despite a very high internal pollution load.*

373 Phytoextraction crop disposal—an unsolved problem
A. SAS-NOWOSIELSKA, R. KUCHARSKI, E. MALKOWSKI, M. POGRZEBA (Poland), J.M. KUPERBERG (USA), K. KRYŃSKI (Poland)
Methods of contaminated crop disposal are described and evaluated.

381 Organochlorine residues in harbour porpoises from Southwest Greenland
A. BORRELL, A. AGUILAR, G. CANTOS (Spain), C. LOCKYER, M.P. HEIDE-JØRGENSEN, J. JENSEN (Denmark)
While organochlorine levels in W. Greenland harbour porpoises are low, their contribution to Inuit dietary intake should not be disregarded.

393 Antioxidant responses to benzo[*a*]pyrene and Aroclor 1254 exposure in the green-lipped mussel, *Perna viridis*
C.C.C. CHEUNG, W.H.L. SIU, B.J. RICHARDSON, S.B. DE LUCA-ABBOTT, P.K.S. LAM (Hong Kong)
Benzo[a]pyrene and Aroclor 1254 separately and in combination caused specific induction in a number of antioxidant parameters in the gill and hepatopancreas of green-lipped mussels.

405 A cumulative ozone uptake–response relationship for the growth of Norway spruce saplings
P.E. KARLSSON, E.L. MEDIN, S. OTTOSSON, G. SELLDÉN, G. WALLIN, H. PLEIJEL, L. SKÄRBY (Sweden)
Growth reductions by Norway spruce saplings correlate with cumulative needle uptake of ozone.

419 The quantification and distribution of pollution Pb at a woodland in rural south central Ontario, Canada
S.A. WATMOUGH, T.C. HUTCHINSON (Canada)
The distribution of pollution lead was determined at a woodland through the use of stable leads isotopes.

429 Priming effects on PAH degradation and ecotoxicity during a phytoremediation experiment
E.J. JONER (France), D. HIRMANN, O.H.J. SZOLAR, D. TODOROVIC (Austria), C. LEYVAL (France), A.P. LOIBNER (Austria)
Priming effects during set-up of bioremediation laboratory experiments may largely surpass treatment effects.

437 Effects of incremental increases in silt load on the cardiovascular performance of riverine and lacustrine rock bass, *Ambloplites rupestris*
C.M. BUNT (Canada), S.J. COOKE, J.F. SCHREER, D.P. PHILIPP (USA)
Cardiovascular compensation in fish exposed to elevated levels of silt.

Short Communication

445 Invariant chlorine isotopic signatures during microbial PCB reductive dechlorination
N.J. DRENZEK, T.I. EGLINTON, C.O. WIRSEN, N.C. STURCHIO, L.J. HERATY, K.R. SOWERS, Q. WU, H.D. MAY, C.M. REDDY (USA)
No chlorine isotopic fractionation was observed for the microbial reductive dechlorination of 2,3,4,5-tetrachlorobiphenyl.

I Contents of Volume 128